

The basic gas laws

How P , V , and T are related


The "pneumatic" era of chemistry began with the discovery of the *vacuum* around 1650 which clearly established that gases are a form of matter. The ease with which gases could be studied soon led to the discovery of numerous empirical (experimentally-discovered) laws that proved fundamental to the later development of chemistry and led indirectly to the atomic view of matter. These laws are so fundamental to all of natural science and engineering that everyone learning these subjects needs to be familiar with them.

1 Pressure-volume relations: Boyle's law

Robert Boyle (1627-91) showed that the volume of air trapped by a liquid in the closed short limb of a J-shaped tube decreased in exact proportion to the pressure produced by the liquid in the long part of the tube. The trapped air acted much like a spring, exerting a force opposing its compression. Boyle called this effect "*the spring of the air*", and published his results in a pamphlet of that title.

The difference between the heights of the two mercury columns gives the pressure (76 cm = 1 atm), and the volume of the air is calculated from the length of the air column and the tubing diameter.

Some of Boyle's actual data are shown in the table.



volume	pressure	$P \times V$
96.0	2.00	192
76.0	2.54	193
46.0	4.20	193
26.0	7.40	193



Boyle's law states that the pressure of a gas, held at a constant temperature, varies inversely with its volume:

It is very important that you understand how these two equations express the green-highlighted statement of the law. If you feel the need to memorize Eqn 1-2, you probably lack this understanding; please study the paragraph that follows.


$$PV = \text{constant} \quad (1-1) \text{ must know!}$$

or, equivalently,

$$P_1V_1 = P_2V_2 \quad (1-2)$$


Boyle's law is a relation of *inverse proportionality*; any change in the pressure is exactly compensated by an opposing change in the volume. As the pressure decreases toward zero, the volume will increase without limit. Conversely, as the pressure is increased, the volume decreases, but can never reach zero. There will be a separate P - V plot for each temperature; a single P - V plot is therefore called an **isotherm**.

Please take the time to understand this kind of plot which governs any relationship of inverse proportionality, and why the curves never reach the x - and y axes. You should be able to sketch out such a plot when given the value of any one (x,y) pair.




Shown here are some isotherms for one mole of an ideal gas at several different temperatures. Each plot has the shape of a *hyperbola*— the locus of all points having the property $xy = a$, where a is a constant. You will see later how the value of this constant ($PV=25$ for the 300K isotherm shown here) is determined.

A related type of plot with which you should be familiar shows the product PV as a function of the pressure. You should understand why this yields a straight line, and how this set of plots relates to the one immediately above.



Problem Example 1



In an industrial process, a gas confined to a volume of 1 L at a pressure of 20 atm is allowed to flow into a 12-L container by opening the valve that connects the two containers. What will be the final pressure of the gas?

Solution: The final volume of the gas is $(1 + 12)\text{L} = 13\text{ L}$. The gas expands in inverse proportion two volumes

$$P_2 = (20\text{ atm}) \times (1\text{ L} \div 13\text{ L}) = 1.5\text{ atm}$$

Note that there is no need to make explicit use of any "formula" in problems of this kind!

2 How the temperature affects the volume: Charles' law



Jacques Charles

All matter expands when heated, but gases are special in that their degree of expansion is independent of their composition. In 1802, the French scientist [Joseph Gay-Lussac](#) (1778-1850) found that if the pressure is held constant, the volume of any gas changes by the same fractional amount (1/273 of its value) for each C° change in temperature.


On learning that another scientist, [Jacques Charles](#), had observed the proportionality between the volume and temperature of a gas 15 years earlier but had never published his findings, Gay-Lussac generously acknowledged this fact. Eventually, the "law of Charles and Gay-Lussac" became commonly known simply as "Charles' law".

The volume of a gas confined against a constant pressure is directly proportional to the absolute temperature.



Joseph Gay-Lussac

A graphical expression of Charles' law can be seen in these plots of the volume of one mole of an ideal gas as a function of its temperature at various constant pressures.



What do these plots show?

The straight-line plots show that the ratio V/T (and thus dV/dT) is a constant at any given pressure. Thus we can express the law algebraically as

$$V/T = \text{constant} \quad \text{or} \quad V_1/T_1 = V_2/T_2$$

(be sure you understand why these two equations are equivalent!)

What is the significance of the extrapolation to zero volume?

If a gas contracts by 1/273 of its volume for each degree of cooling, it should contract to zero volume at a temperature of -273°C. This, of course, is the [absolute zero of temperature](#), and this extrapolation of Charles' law is the first evidence of the special significance of this temperature.

Why do the plots for different pressures have different slopes?

The lower the pressure, the greater the volume (Boyle's law), so at low pressures the fraction ($V/273$) will have a larger value. You might say that the gas must "contract faster" to reach zero volume when its starting volume is larger.

- Try this [interactive exploration](#) of the Law of Charles and Gay-Lussac
- Carleton University [lecture demonstration video of Charles' Law](#) (YouTube)

Problem Example 2

The air pressure in a car tire is 30 psi (pounds per square inch) at 10°C. What will be pressure be after driving has raised its temperature to 45°C ? (Assume that the volume remains unchanged.)

Solution: The gas expands in direct proportion to the ratio of the absolute temperatures:

$$P_2 = (30 \text{ psi}) \times (318\text{K} \div 283\text{K}) = 33.7 \text{ psi}$$

Historical notes

The fact that a gas expands when it is heated has long been known. In 1702, [Guillaume Amontons](#) (1163-1705), who is better known for his early studies of friction, devised a thermometer that related the temperature to the volume of a gas. Robert Boyle had observed this relationship in 1662, but the lack of any uniform temperature scale at the time prevented them from establishing the relationship as we presently understand it.

Jacques Charles discovered the law that is named for him in the 1780s, but did not publish his work. John Dalton published a form of the law in 1801, but the first thorough published presentation was made by Gay-Lussac in 1802, who acknowledged Charles' earlier studies.

Hot-air balloons and Charles' Law



The buoyancy that lifts a hot-air balloon into the sky depends on the difference between the density (mass ÷ volume) of the air entrapped within the balloon's envelope, compared to that of the air surrounding it. When a balloon on the ground is being prepared for flight, it is first partially inflated by an external fan, and possesses no buoyancy at all. Once the propane burners are started, this air begins to expand according to Charles' law. After the warmed air has completely inflated the balloon, further expansion simply forces excess air out of the balloon, leaving the weight of the diminished mass of air inside the envelope smaller than that of the greater mass of cooler air that the balloon displaces.




Jacques Charles collaborated with the Montgolfier Brothers whose hot-air balloon made the world's first manned balloon flight in June, 1783 (left). Ten days later, Charles himself co-piloted the first hydrogen-filled balloon. Gay-Lussac (right), who had a special interest in the composition of the atmosphere, also saw the potential of the hot-air balloon, and in 1804 he ascended to a then-record height of 6.4 km. [images: Wikimedia]



If you ever visit Paris, look for a street and a hotel near the Sorbonne that bear Gay-Lussac's name.

Jacques Charles' understanding of buoyancy led to his early interest in balloons and to the use of hydrogen as an inflating gas. His first flight was witnessed by a crowd of 40,000 which included Benjamin Franklin, at that time ambassador to France. On its landing in the countryside, the balloon was reportedly attacked with axes and pitchforks by terrified peasants who believed it to be a monster from the skies. Charles' work was mostly in mathematics, but he managed to invent a number of scientific instruments and confirmed experiments in electricity that had been performed earlier by Benjamin Franklin and others.[*]

Gay-Lussac's balloon flights enabled him to sample the composition of the atmosphere at different altitudes (he found no difference). His Law of Combining Volumes (described below) constituted one of the foundations of modern chemistry. Gay-Lussac's contributions to chemistry are numerous; his work in electrochemistry enabled him to produce large quantities of sodium and potassium; the availability of these highly active metals led to his co-discovery of the element boron. He was also the first to recognize iodine as an element. In an entirely different area, he developed a practical method of measuring the alcohol content of beverages, and coined the names "pipette" and "burette" that are known to all chemistry students. A good biography can be viewed at Google Books.



3 Volume and the number of molecules: Avogadro's law

Gay-Lussac's Law of Combining Volumes

For a good tutorial overview of the Law of Combining Volumes, see this [ChemPaths page](#).

In the same 1808 article in which Gay-Lussac published his observations on the thermal expansion of gases, he pointed out that **when two gases react, they do so in volume ratios that can always be expressed as small whole numbers**. This came to be

known as the **Law of combining volumes**.

These "small whole numbers" are of course the same ones that describe the "combining weights" of elements to form simple compounds, as described in the lesson dealing with [simplest formulas from experimental data](#).

The Italian scientist [Amedeo Avogadro](#) (1776-1856) drew the crucial conclusion: these volume ratios must be related to the relative numbers of molecules that react, and thus the famous "E.V.E.N principle":

Equal **v**olumes of gases, measured at the same temperature and pressure, contain **e**qual **n**umbers of molecules

Avogadro's law thus predicts a *directly proportional* relation between the number of moles of a gas and its volume.

This relationship, originally known as *Avogadro's Hypothesis*, was crucial in establishing the formulas of simple molecules at a time (around 1811) when the distinction between atoms and molecules was not clearly understood. In particular, the existence of diatomic molecules of elements such as H_2 , O_2 , and Cl_2 was not recognized until the results of combining-volume experiments such as those depicted below could be interpreted in terms of



Amedeo Avogadro


the E.V.E.N. principle.

How the E.V.E.N. principle led to the correct formula of water


Early chemists made the mistake of assuming that the formula of water is HO. This led them to miscalculate the molecular weight of oxygen as 8 (instead of 16). If this were true, the reaction $\text{H} + \text{O} \rightarrow \text{HO}$ would correspond to the following combining volumes results according to the E.V.E.N principle:



But a similar experiment on the formation of hydrogen chloride from hydrogen and chlorine yielded twice the volume of HCl that was predicted by the the assumed reaction $\text{H} + \text{Cl} \rightarrow \text{HCl}$. This could be explained only if hydrogen and chlorine were diatomic molecules:




This made it necessary to re-visit the question of the formula of water. The experiment immediately confirmed that the correct formula of water is H_2O :



This conclusion was also seen to be consistent with the observation, made a few years earlier by the English chemists Nicholson and Carlisle that the reverse of the above reaction, brought about by the electrolytic decomposition of water, yields hydrogen and oxygen in a 2:1 volume ratio.

A nice overview of these developments can be seen at [David Dice's Chemistry page](#), from which this illustration is taken.



4 The ideal gas equation of state

If the variables P , V , T and n (the number of moles) have known values, then a gas is said to be in a definite **state**, meaning that all other physical properties of the gas are also defined. The relation between these *state variables* is known as an **equation of state**. By combining the expressions of Boyle's, Charles', and Avogadro's laws (you should be able to do this!) we can write the very important **ideal gas** equation of state

$$PV = nRT$$

Must know this!

where the proportionality constant R is known as the **gas constant**. This is one of the few equations you *must* commit to memory in this course; you should also know the common value and units of R .


An *ideal gas* is defined as a hypothetical substance that obeys the ideal gas equation of state.

Take note of the word "hypothetical" here. No real gas (whose molecules occupy space and interact with each other) can behave in a truly ideal manner. But we will see in the [last lesson](#) of this series that all *gases* behave more and more like an ideal gas as the pressure approaches zero. A pressure of only 1 atm is sufficiently close to zero to make this relation useful for most gases at this pressure.

Many textbooks show formulas, such as $P_1V_1 = P_2V_2$ for Boyle's law. *Don't bother memorizing them*; if you really understand the meanings of these laws as stated above, you can easily derive them on the rare occasions when they are needed. **The ideal gas equation is the only one you need to know.**

***PVT* surface for an ideal gas**

In order to depict the relations between the three variables P , V and T we need a three-dimensional graph.



Each point on the curved surface represents a possible combination of (P, V, T) for an arbitrary quantity of an ideal gas. The three sets of lines inscribed on the surface correspond to states in which one of these three variables is held constant.

The red curved lines, being lines of constant temperature, or *isotherms*, are plots of Boyle's law. These isotherms are also seen projected onto the P - V plane at the top right.


The yellow lines are *isochors* and represent changes of the pressure with temperature at constant volume.

The green lines, known as *isobars*, and projected onto the V - T plane at the bottom, show how the volumes contract to zero as the absolute temperature approaches zero, in accordance with the law of Charles and Gay-Lussac.

Problem Example 3

A biscuit made with baking powder has a volume of 20 mL, of which one-fourth consists of empty space created by gas bubbles produced when the baking powder decomposed to CO₂. What weight of NaHCO₃ was present in the baking powder in the biscuit? Assume that the gas reached its final volume during the baking process when the temperature was 400°C.

(Baking powder consists of sodium bicarbonate mixed with some other solid that produces an acidic solution on addition of water, initiating the reaction



Solution: Use the ideal gas equation to find the number of moles of CO₂ gas; this will be the same as the number of moles of NaHCO₃ (84 g mol⁻¹) consumed :

$$n = \frac{(1 \text{ atm}) (0.005 \text{ L})}{(.082 \text{ L atm mol}^{-1} \text{ K}^{-1})(673\text{K})} = 9.1\text{E-}6 \text{ mol}$$


$$9.1\text{E-}6 \text{ mol} \times 84 \text{ g mol}^{-1} = 0.0076 \text{ g}$$

What you should be able to do

Make sure you thoroughly understand the following essential ideas which have been presented above, and be able to state them in your own words. It is especially important that you know the precise meanings of all the green-highlighted terms in the context of this topic.

- **Boyle's Law** - The *PV* product for any gas at a fixed temperature has a constant value. Understand how this implies an *inverse* relationship between the pressure and the volume.
- **Charles' Law** - The volume of a gas confined by a fixed pressure varies *directly* with the absolute temperature. The same is true of the pressure of a gas confined to a fixed volume.
- **Avogadro's Law** - This is quite intuitive: the volume of a gas confined by a fixed pressure varies directly with the quantity of gas.
- The E.V.E.N. principle - this is just another way of expressing Avogadro's Law.
- Gay-Lussac's Law of Combining Volumes - you should be able to explain how this principle, that follows from the E.V.E.N. principle and the Law of Combining Weights,
- **The ideal gas equation of state** - this is one of the very few mathematical relations you *must* know. Not only does it define the properties of the hypothetical substance known as an **ideal gas**, but its importance extends quite beyond the subject of gases.

Concept Map



© 2004-2017 by Stephen Lower - last modified 2017-08-15

The Chem1 Virtual Textbook home page is at <http://www.chem1.com/acad/virtualtextbook.html>



This work is licensed under a Creative Commons Attribution 3.0 Unported License